

REPORT DOCUMENTATION PAGE		Form Approved OMB NO. 0704-0188	
Public Reporting Burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comment regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA, 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington DC 20503			
1. AGENCY USE ONLY (Leave Blank)		2. REPORT DATE:	
		3. REPORT TYPE AND DATES COVERED Final Report 10-Jul-2003 - 9-Jul-2006	
4. TITLE AND SUBTITLE Modeling Complex Nonlinear Optical Systems		5. FUNDING NUMBERS DAAD190310209	
6. AUTHORS Alejandro Aceves, Todd Kapitula		8. PERFORMING ORGANIZATION REPORT NUMBER	
7. PERFORMING ORGANIZATION NAMES AND ADDRESSES University of New Mexico Contracts & Grants, Office Of Research Services 1 University of New Mexico Albuquerque, NM 87131 -0001			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709-2211		10. SPONSORING / MONITORING AGENCY REPORT NUMBER 45248-PH-HSI.1	
11. SUPPLEMENTARY NOTES The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision, unless so designated by other documentation.			
12. DISTRIBUTION AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The abstract is below since many authors do not follow the 200 word limit			
14. SUBJECT TERMS Nonlinear Optics, Photonics, Bose Einstein Condensates, Applied Mathematics		15. NUMBER OF PAGES Unknown due to possible attachments	
		16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION ON THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL

Report Title

Modeling Complex Nonlinear Optical Systems

ABSTRACT

This research dealt with the modeling of light propagation in nonlinear periodic media including bragg grating fiber arrays and periodic nonlinear 2-dimensional waveguides. The goals set were to find conditions for stable pulse propagation in the arrays and for the search of light bullets, their stability and propagation characteristics in the two dimensional waveguide. We also established conditions for optical trapping in a defect. This is a topic of great interest in the search of all optical logic systems and buffers. A second component of the project dealt with existence and stability of Bose Einstein condensates in periodic magnetic traps. There has been an extensive experimental effort on BEC trapping and our work developed a solid theoretical framework to explore such trapping mechanisms. Tools used in this research include: Dynamical systems, numerical methods for nonlinear partial differential equations, asymptotic analysis. The project had also an important educational component as it served to train 3 graduate students in Applied Mathematics and provided a seed for a new crop of students working in this field.

List of papers submitted or published that acknowledge ARO support during this reporting period. List the papers, including journal references, in the following categories:

(a) Papers published in peer-reviewed journals (N/A for none)

T. Dohnal, A. Aceves (2005), "Optical soliton bullets in (2+1)D nonlinear bragg resonant periodic geometries", *Studies in Applied Mathematics*, **115** 209-232.

T. Dohnal and A. B. Aceves (2006) "Finite dimensional model for defect-trapped light in planar periodic nonlinear structures", *Optics Letters* **31**, 3013-3015.

T. Dohnal and T. Hagstrom (2006), "Perfectly matched layers in photonics computations: 1D and 2D Nonlinear Coupled Mode Equations," accepted to *J. Comput. Phys.*

T. Kapitula, B. Sandstede (2004), "Eigenvalues and resonances using the Evans function", *Discrete and Continuous Dynamical Systems*, Vol. 10, No. 4, 857-869.

T. Kapitula, B. Sandstede and J. Kutz (2004), "The Evans function for nonlocal equations", *Indiana U. Math. J.*, Vol. 53, No. 4, 1095-1126.

T. Kapitula and P. Kevrekidis (2004), "Linear stability of perturbed Hamiltonian systems: theory and a case example", *J. Phys. A: Math. Gen.*, Vol. 37, No. 30, 7509-7526.

T. Kapitula, P. Kevrekidis and B. Sandstede (2004), "Counting eigenvalues via the Krein signature in infinite-dimensional Hamiltonian systems" *Physica D*, Vol. 195, No. 3&4, 263-282.
(Addendum: Counting eigenvalues via the Krein signature in infinite-dimensional Hamiltonian systems", *Physica D*, Vol. 201, No. 1&2, 199-201 (2005).

T. Kapitula (2005), "Stability analysis of pulses via the Evans function: dissipative systems", *Lecture Notes in Physics*, Vol. 661, 407-427.

T. Kapitula and P. Kevrekidis (2005), "Bose-Einstein condensates in the presence of a magnetic trap and optical lattice: two-mode approximation", *Nonlinearity*, Vol. 18, No. 6, 2491-2512.

T. Kapitula and P. Kevrekidis (2005), "Bose-Einstein condensates in the presence of a magnetic trap and optical lattice", *Chaos* Vol. 15, No. 3, 037114.

Number of Papers published in peer-reviewed journals: 10.00

(b) Papers published in non-peer-reviewed journals or in conference proceedings (N/A for none)

A.B. Aceves and T. Dohnal, "Stopping and bending light in 2D photonic structures." "Nonlinear Waves: Classical and Quantum Effects," p. 293 - 302, F. Kh. Abdullaev and V.V. Konotop (eds.), Kluwer, (2004).

A.B. Aceves and T. Dohnal, "Stopping and bending light in 2D photonic structures." *Proceedings of OSA topical meeting on Nonlinear Guided Waves and their Applications*, Toronto, March (2004).

Number of Papers published in non peer-reviewed journals: 2.00

(c) Presentations

Number of Presentations: 0.00

Non Peer-Reviewed Conference Proceeding publications (other than abstracts):

Number of Non Peer-Reviewed Conference Proceeding publications (other than abstracts): 0

Peer-Reviewed Conference Proceeding publications (other than abstracts):

Number of Peer-Reviewed Conference Proceeding publications (other than abstracts): 0

(d) Manuscripts

"Three is a crowd: Solitary waves in photorefractive media with three potential wells", T. Kapitula, P. Kevrekidis and Z. Chen, to appear in SIAM J. Appl. Dyn. Sys.

Number of Manuscripts: 0.00

Number of Inventions:

Graduate Students

<u>NAME</u>	<u>PERCENT SUPPORTED</u>	
Tomas Dohnal	1.00	No
Karl Frinkle	0.75	No
Bobbi Page	0.25	No
FTE Equivalent:	2.00	
Total Number:	3	

Names of Post Doctorates

<u>NAME</u>	<u>PERCENT SUPPORTED</u>
FTE Equivalent:	
Total Number:	

Names of Faculty Supported

<u>NAME</u>	<u>PERCENT SUPPORTED</u>	National Academy Member
Alejandro Aceves	0.50	No
Todd Kapitula	0.50	No
FTE Equivalent:	1.00	
Total Number:	2	

Names of Under Graduate students supported

<u>NAME</u>	<u>PERCENT SUPPORTED</u>
FTE Equivalent:	
Total Number:	

Names of Personnel receiving masters degrees

<u>NAME</u>	
Bobbi Page	No
Total Number:	1

Names of personnel receiving PHDs

<u>NAME</u>	
Tomas Dohnal	No
Karl Frinkle	No
Total Number:	2

Names of other research staff

<u>NAME</u>	<u>PERCENT SUPPORTED</u>
FTE Equivalent:	
Total Number:	

Sub Contractors (DD882)

Inventions (DD882)




The University of New Mexico

Modeling Complex Nonlinear Optical Systems

Alejandro Aceves

Department of Mathematics and Statistics

The University of New Mexico



Statement of the problem:

Study the dynamics of pulses in photonic structures.

What is a photonic structure ?

- It is an “engineered” optical medium with periodic properties.
- Photonic structures are built to manipulate light (slow light, light localization, trap light)
- I’ll show some examples, but this talk will concentrate on 2-dim periodic waveguides



Motivation of study

We take advantage of periodic structures and nonlinear effects to propose new stable and robust systems relevant to optical systems.

- Periodic structure - material with a periodically varying index of refraction (“grating”)
- Nonlinearity - dependence of the refractive index on the intensity of the electric field (Kerr nonlinearity)


The effects we seek:

- existence of stable localized solutions – solitary waves, solitons
- short formation lengths of these stable pulses
- possibility to control the pulses – speed, direction (2D, 3D)

Prospective applications:

- rerouting of pulses
- optical memory
- low-loss bending of light

1d structures: Optical fiber gratings



$E_f(Z, T)$ (forward moving envelope)

$E_b(Z, T)$ (backward moving envelope)



Equations studied

1-dim Coupled Mode Equations

$$\partial_t E_+ = -c_g \partial_z E_+ + i\kappa E_- + i\Gamma(|E_+|^2 + 2|E_-|^2)E_+$$


$$\partial_t E_- = c_g \partial_z E_- + i\kappa E_+ + i\Gamma(|E_-|^2 + 2|E_+|^2)E_-$$

Gap solitons exist. Velocity proportional to amplitude mismatch
Between forward and backward envelopes.

II. 2D structures: Waveguide gratings

Assumptions:

- dynamics in y arrested by a fixed $n(y)$ profile
- xy -normal incidence of pulses
- characteristic length scales of coupling, nonlinearity and diffraction are in balance



BARE 2D WAVEGUIDE


- 2D NL Schrödinger equation
- collapse phenomena: point blow-up

WAVEGUIDE GRATING

- 2D CME
- no collapse
- possibility of localization



Dispersion relation for coupled mode equations



Governing equations : 2D Coupled Mode Equations

$$\partial_t E_+ = -c_g \partial_z E_+ + id \partial_{x^2} E_+ + i\kappa E_- + i\Gamma(|E_+|^2 + 2|E_-|^2)E_+$$

$$\partial_t E_- = c_g \partial_z E_- + id \partial_{x^2} E_- + i\kappa E_+ + i\Gamma(|E_-|^2 + 2|E_+|^2)E_-$$

advection

diffraction

coupling

non-linearity

$$c_g, d, \kappa, \Gamma \geq 0, \quad E_{\pm} : [-L_x, L_x] \times [-L_z, L_z] \times [0, \infty) \rightarrow C.$$

Summary of results (details to follow on next slides)

- **Found stationary solutions of the governing equations of the 2d structure (see next slide)**
- **Obtained conditions for bullet propagation in such structures**
- **Obtained conditions for light trapping at a defect by a resonance mechanism between the incident optical bullets and defect modes**
- **Derived a finite dimensional dynamical system to study the dynamics inside the trap**

Stationary solutions via Newton's iteration

If $E_{\pm}(x, z, t) = \mathcal{E}_{\pm}(x, z) e^{-i\omega t}$ then

$$\begin{aligned}\omega \mathcal{E}_+ + i c_g \partial_z \mathcal{E}_+ + \partial_x^2 \mathcal{E}_+ + \kappa \mathcal{E}_- + \Gamma(|\mathcal{E}_+|^2 + 2|\mathcal{E}_-|^2) \mathcal{E}_+ &= 0, \\ \omega \mathcal{E}_- - i c_g \partial_z \mathcal{E}_- + \partial_x^2 \mathcal{E}_- + \kappa \mathcal{E}_+ + \Gamma(|\mathcal{E}_-|^2 + 2|\mathcal{E}_+|^2) \mathcal{E}_- &= 0.\end{aligned}\tag{1}$$

Solve (1) as a **NL eigenvalue problem** for $\left(\omega, \begin{pmatrix} \mathcal{E}_+ \\ \mathcal{E}_- \end{pmatrix}\right)$ via Newton's iteration.

Need one more equation:

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |\mathcal{E}_+|^2 + |\mathcal{E}_-|^2 dz dx = N.$$

Initial guess: $\left(\omega^{(0)}, \mathcal{E}_{\pm}^{(0)}(x, z)\right)$

separable waveform $\mathcal{E}_{\pm}^{(0)}(x, z) = \mathcal{F}_{\pm}(z) G(x), \quad \omega^{(0)} = \kappa \cos(\delta)$


where $\mathcal{F}_{\pm}(z) e^{-i\omega^{(0)}t}$ is the 1D gap soliton with $v = 0$ (free parameter $\delta \in (0, \pi)$)

Substitute and integrate in z :

$$G'' + b(G^3 - G) = 0, \quad b = 2 \frac{\kappa}{\delta} (\sin(\delta) - \delta \cos(\delta)), \quad \delta \in (0, \pi/2)$$

$$\Rightarrow G(x) = \sqrt{2} \operatorname{sech}(\sqrt{b}x)$$

Stationary case I



Nonexistence of minima of the Hamiltonian

Theorem 1. *The Hamiltonian functional of 2D CME has no minima constrained to a fixed total power.*

Proof: Suppose $\exists(\mathcal{E}_+(x, z), \mathcal{E}_-(x, z))$ s.t. $H(\mathcal{E}_+, \mathcal{E}_-) = \min_S H$, where

$$S = \{(f_1(x, z), f_2(x, z)) \text{ s.t. } f_{1,2} : \mathbb{R}^2 \rightarrow \mathbb{C}, \sum_{k=1}^2 \|f_k\|_2^2 = \|\mathcal{E}_+\|_2^2 + \|\mathcal{E}_-\|_2^2\}$$

Consider

$$S_1 = \{(\tilde{\mathcal{E}}_+, \tilde{\mathcal{E}}_-) : \tilde{\mathcal{E}}_{\pm} = \alpha \mathcal{E}_{\pm}(x/\mu, z/\nu) \text{ and } \alpha, \mu, \nu > 0, \alpha^2 \mu \nu = 1\}.$$

Clearly $S_1 \subset S$ and $(\mathcal{E}_+, \mathcal{E}_-) \in S_1$.

Within S_1 the Hamiltonian $H = H_r = A_1 \frac{1}{\nu} + A_3 \alpha^4 \nu^2 - A_4 \alpha^2 - A_2$ with

$$A_1 = ic_g \int_{\mathbb{R}^2} \mathcal{E}_-^* \partial_z \mathcal{E}_- - \mathcal{E}_+^* \partial_z \mathcal{E}_+ dx dz, \quad A_2 = \kappa \int_{\mathbb{R}^2} \mathcal{E}_- \mathcal{E}_+^* + \mathcal{E}_-^* \mathcal{E}_+ dx dz,$$

$$A_3 = \int_{\mathbb{R}^2} |\partial_x \mathcal{E}_+|^2 + |\partial_x \mathcal{E}_-|^2 dx dz, \quad A_4 = \frac{\Gamma}{2} \int_{\mathbb{R}^2} |\mathcal{E}_+|^4 + 4|\mathcal{E}_+|^2 |\mathcal{E}_-|^2 + |\mathcal{E}_-|^4 dx dz$$

$A_1, A_2 \in \mathbb{R}$ and $A_3, A_4 > 0$. The only C.P. of H_r is


$$(\alpha^*, \nu^*) = \left(\frac{A_1 \sqrt{2A_3}}{A_4^{3/2}}, \frac{A_4^2}{2A_1 A_3} \right)$$

and by the 2nd derivative test (α^*, ν^*) is a **saddle**!

□


Defects in 1d gratings

- Trapping of a gap soliton bullet in a defect. (this concept has been theoretically demonstrated by Goodman, Weinstein, Slusher for the 1-dim (fiber Bragg grating with a defect) case).



Ref: R. Goodman et.al., JOSA B 19, 1635 (July 2002)

Governing equations : 2D CME



$$\begin{aligned}\partial_t E_+ &= -\partial_z E_+ + i\partial_{x^2} E_+ + i\kappa(x, z)E_- + V(x, z)E_+ + i\Gamma(|E_+|^2 + 2|E_-|^2)E_+ \\ \partial_t E_- &= \partial_z E_- + i\partial_{x^2} E_- + i\kappa(x, z)E_+ + V(x, z)E_- + i\Gamma(|E_-|^2 + 2|E_+|^2)E_-\end{aligned}$$

advection

diffraction


coupling

defect

non-linearity


$$\kappa, \Gamma \geq 0, \quad E_{\pm} : [-L_x, L_x] \times [-L_z, L_z] \times [0, \infty) \rightarrow \mathbb{C}.$$

2-D version of resonant trapping



left: GS modulus

right: defect potential $V(x, z)$



(-) Bifurcation curves for the NL defect mod

(*) stationary GS with $\omega_0 \approx 0.96$

3 trapping cases (GS: $\omega(v=0) \approx 0.96$)

1. Trapping into **two** defect modes


$$V = V_1(x)T(z;9) + V_2(z)T(x;7), \quad \kappa = \sqrt{1 + k^2(\tanh^2(kz) - 1)},$$

$$\text{where } V_1 = 2\beta^2 \text{sech}^2(\beta x), \quad V_2 = \frac{1}{2} \frac{k^2 \sqrt{1 - k^2} \text{sech}^2(kz)}{1 + k^2(\tanh^2(kz) - 1)} \quad \text{and} \quad T(y; c) = \frac{1}{2}(\tanh(y + c) - \tanh(y - c))$$


with $k = 0.18$ and $\beta = 0.16$

\Rightarrow 2 linear defect modes: $\omega_L \approx 0.963, 0.992$

$$|E_+(x=0, z, t)|$$



$$|a_k(t)|$$




3 trapping cases (GS: $\omega(v=0) \approx 0.96$)

2. Trapping into **one** defect mode

$$V = 0.3e^{-(ax^2+bz^2)}, \quad \kappa = 1 + 0.1e^{-(ax^2+bz^2)}, \quad a = 0.25, b = 0.3$$

\Rightarrow 1 linear defect mode : $\omega_L \approx 0.995$



3 trapping cases (GS: $\omega(v=0) \approx 0.96$)



3. No trapping

$$V = V_1(x)T(z;5) + V_2(z)T(x;5), \quad \kappa = \sqrt{1 + k^2(\tanh^2(kz) - 1)},$$

$$\text{where } V_1 = 2\beta^2 \text{sech}^2(\beta x), \quad V_2 = \frac{1}{2} \frac{k^2 \sqrt{1 - k^2} \text{sech}^2(kz)}{1 + k^2(\tanh^2(kz) - 1)} \quad \text{and} \quad T(y; c) = \frac{1}{2}(\tanh(y + c) - \tanh(y - c))$$

with $k = 0.85$ and $\beta = 1$

\Rightarrow 5 linear defect modes: $\omega_L \approx -0.47, 0.26, 0.47, 0.68$ and 0.87 ← ALL FAR FROM RESONANCE



Finite dimensional approximation of the trapped dynamics

For trapped solutions with small amplitude:

$$\begin{pmatrix} E_+(x, z, t) \\ E_-(x, z, t) \end{pmatrix} \approx \sum_{k=1}^N a_k(t) e^{-i\omega_{L_k} t} \begin{pmatrix} \psi_{+k}(x, z) \\ \psi_{-k}(x, z) \end{pmatrix},$$

where $(\psi_{+k}, \psi_{-k})^T, k = 1, \dots, N$ are the defect modes.

Substituting into 2D CME with the defect potentials

$$i \sum_{k=1}^N a'_k \begin{pmatrix} \psi_{+k} \\ \psi_{-k} \end{pmatrix} + \Gamma \begin{pmatrix} (NL)_+ \\ (NL)_- \end{pmatrix} = 0,$$

where $(NL)_\pm = (|E_\pm|^2 + 2|E_\mp|^2)E_\pm$.

Due to orthogonality

$$ia'_k(t) + \Gamma \int (NL)_+ \psi_{+k}^* + (NL)_- \psi_{-k}^* dx dz = 0, \quad k = 1, \dots, N.$$


Finite dimensional approximation of the trapped dynamics

For $N = 2$:

after transformation $\tilde{a}_{1,2}(t) = a_{1,2}(t)e^{\mp \frac{i\Delta\omega}{2}t}$, where $\Delta\omega = \omega_{L_1} - \omega_{L_2}$

$$i\tilde{a}'_1 - \frac{\Delta\omega}{2}\tilde{a}_1 + \alpha_1|\tilde{a}_1|^2\tilde{a}_1 + \beta|\tilde{a}_2|^2\tilde{a}_1 + \frac{\beta}{2}\tilde{a}_2^2\tilde{a}_1^* = 0$$

$$i\tilde{a}'_2 + \frac{\Delta\omega}{2}\tilde{a}_2 + \alpha_2|\tilde{a}_2|^2\tilde{a}_2 + \beta|\tilde{a}_1|^2\tilde{a}_2 + \frac{\beta}{2}\tilde{a}_1^2\tilde{a}_2^* = 0.$$





Conclusions (Part I)

- 2d nonlinear photonic structures show promising properties for quasi-stable propagation of “slow” light-bullets
- With the addition of defects, we presented examples of resonant trapping
- Research in line with other interesting schemes to slow down light for eventually having all optical logic devices (eg. buffers)



Part II: Dynamics in Bose-Einstein Condensates

Statement of the problem:

In the study of the dynamics of matter waves for Bose-Einstein condensates, it is of great importance to understand the scenarios under which solutions such as necklaces, multi-poles, and vortices will exist and persist.



Part 2: Bose Einstein Condensates (summary of most important results)

- Shown that, under suitable assumptions, that N-solitons are stable for a large class of integrable partial differential equations including the equation that govern BEC
- Shown how the presence of an optical lattice along with the magnetic trap can influence the dynamics of matter waves in Bose-Einstein condensates
- Illustrated the manner in which increasing the number of potential wells in photorefractive media effects the dynamics associated with solitary waves
- Begun to develop some ``rules-of-thumb" regarding the existence and stability of waves in two-dimensional Bose-Einstein condensates



Bibliography

- “Bose-Einstein condensates in the presence of a magnetic trap and optical lattice: two-mode approximation, T. Kapitula, P. Kevrekidis, Nonlinearity, Vol. 18, No. 6, 2491-2512 (2005)
- “Bose-Einstein condensates in the presence of a magnetic trap and optical lattice”, T. Kapitula, P. Kevrekidis, Chaos Vol. 15, No. 3, 037114 (2005)
- R.E. Slusher and B.J. Eggleton Eds., “Nonlinear Photonic Crystals,” (Springer-Verlag, Berlin, 2003)
- “Optical soliton bullets in (2+1)D nonlinear bragg resonant periodic geometries”, Studies in Applied Mathematics, 115, 209-232, T. Dohnal, A. Aceves (2005).
- “Finite-dimensional model for defect-trapped light in planar periodic nonlinear structures”, Alejandro B. Aceves and Tomáš Dohnal Optics Letters, Vol. 31, Issue 20, pp. 3013-3015 (2006).